

# Injuries of the Hip

Physiotherapy & Occupational Therapy  
01902 307999 ext. 88536

## Introduction

The Doctor has diagnosed you as having a fracture or a soft tissue injury. This information leaflet has been written to help you manage this injury at home. An appointment may also be arranged for you to see a Physiotherapist.

## Have you been told you have a fracture?

A fracture (or a break) is damage to a bone. This may be a small chip of bone or a complete break through the bone.

## How are fractures treated?

Fractures may be treated in a number of ways:

- Plaster cast
- Internal fixation is when the fracture is fixed with nails, plates, screws, or wires
- External fixation is when nails are placed into the bone with a metal plate outside to hold the bone still while the fracture heals
- Reduction in theatre – the bone is put into a good position. This is often followed by one of the above treatments.

Some fractures are not treated in any of the ways outlined above. They may be managed with a tubular bandage, brace, or splint and you may be encouraged to begin exercising as soon as possible.

## Have you been told you have a soft tissue injury?

This includes sprains, strains, and bruises which affect the soft tissues of the body. Most commonly injured are the ligaments, muscles, tendons, and joint capsules. Following a fracture the soft tissues will also have been injured.

There is no set time for a fracture or a soft tissue injury to heal. The Doctors, Nurses, and Physiotherapists will give you advice on what you can expect and what you should be doing. You should expect some swelling and discomfort for several weeks, particularly after you have been on your feet for long periods.

**The prevention of infection is a major priority in all healthcare and everyone has a part to play.**

- Please decontaminate your hands frequently for 20 seconds using soap and water or alcohol gel if available
- If you have symptoms of diarrhoea and/or vomiting, cough or other respiratory symptoms, a temperature or any loss of taste or smell please do not visit the hospital or any other care facility and seek advice from 111
- Keep the environment clean and tidy
- Let's work together to keep infections out of our hospitals and care homes.

## How can I manage my injury at home?

The regime of protection, rest, compression, and elevation is a simple of effective method of early treatment, particularly when swelling is present.

- **Protection** – this may include walking aids, a brace or splint, or tubular bandage
- **Relative rest** – rest the affected area for the first few days, avoiding excessive activity, using any of the above supports you have been given
- **Compression** – this is used to reduce swelling. If you have been given a tubular bandage, wear this whenever the injured area is not elevated and always remove it before you go to bed
- **Elevation** – this helps to reduce swelling and inflammation. Elevate the injured area as much as possible in the first few days, this should be done for at least 20 minutes every two hours.

**Do not use compression when elevating the injured area.**

**For the first three days DO NOT apply heat, massage the injury, or drink alcohol as these activities may increase bleeding or swelling.**

## Do I need a walking aid?

The Physiotherapist will advise you on the most appropriate walking aid and teach you how to use it. Unless otherwise instructed and as you feel able, gradually increase the amount of weight you put through your leg, until you no longer need the walking aid.

**When you no longer need the walking aid please return it to Fracture Clinic.**

## Should I take pain relief?

Your Doctor or Pharmacist can advise on appropriate pain relief which, if taken regularly, can improve comfort and reduce muscle spasms. If your pain is well controlled you will manage exercises more easily which will aid recovery.

## Can I drive?

Your Doctor or Physiotherapist will advise you when it is safe to return to driving. You must be in full control of the vehicle before attempting to drive again. You should not drive whilst wearing a brace or splint.

## Why exercise?

Early exercise can have a positive influence on tissue healing and reduce the potential for joint and soft tissue stiffness. Too much movement or the wrong type of exercise however may be harmful to tissue healing and it may cause inflammation which can increase your pain. If you are not sure if it is safe to do something please talk to your physiotherapist first.

## Exercises – to be completed 3-5 times a day

Lying or sitting. Bend and straighten your leg.  
Repeat 5-10 times.



Lying on your back. Bring your leg out to the side and then back to mid position. Repeat 5-10 times.



Lying or sitting with legs straight. Tighten your thigh muscle to push the knee straight.  
Hold 5 seconds. Repeat 10 times.



Lying or sitting. Bend the unaffected leg and put your foot on the bed. Put a rolled towel under the affected knee. Exercise your affected leg by pulling your foot and toes up, tightening your thigh muscle and straightening the knee (keep knee on the towel). Hold 5 seconds. Repeat 5-10 times.



Lying on your back with knees bent. Squeeze your buttocks together and lift your bottom off the floor. Hold for 5 seconds. Return to starting position. Repeat 5-10 times.



## Other exercises to progress to

Sitting with your arms crossed. Stand up and then sit down slowly on a chair. (This can be made easier and more difficult by changing the height of the chair). Repeat 10 times.



Stand straight holding on to a table or work surface. Bring your leg backwards keeping your knee straight. Do not lean forwards. Repeat 10 times.



Stand straight holding on to a support. Lift your leg sideways and bring it back keeping your body straight throughout the exercise. Repeat 10 times.



Stand straight with support if needed. Lift your affected knee. Repeat 10 times.



Stand in front of a step (with support if required). Step up with the injured leg leading and back down with the uninjured leg. Repeat 10 times.



Lie on your side with your knees bent. Tighten your buttocks. Lift your top knee as far as you can, without letting your pelvis rotate forward or back. Keep your feet together and back straight during the exercise. Repeat 10 times.



## Gradually

- Return to your normal activities
- Increase your walking distance
- Resume hobbies and sports.

If you are unsure of when to progress your exercises, or you have any concerns about your injury and you do not have a further appointment with the physiotherapy team, please contact the physiotherapy department on the telephone number on the front of this leaflet.

## Contact details

The team is available Monday to Friday, 8.00 16.00. Outside of these hours please contact the Emergency Department or, if you have a fracture, the Fracture Clinic. For both of these options please ring 01902 307999 and ask the operator to put you through to the appropriate department.

### **Acute Trauma Physiotherapy Team**

Emergency Department

New Cross Hospital

01902 307999 ext. 88536

## **English**

**If you need information in another way like easy read or a different language please let us know.**

**If you need an interpreter or assistance please let us know.**

## **Lithuanian**

**Jeigu norėtumėte, kad informacija jums būtų pateikta kitu būdu, pavyzdžiui, supaprastinta forma ar kita kalba, prašome mums apie tai pranešti.**

**Jeigu jums reikia vertėjo ar kitos pagalbos, prašome mums apie tai pranešti.**

## **Polish**

**Jeżeli chcieliby Państwo otrzymać te informacje w innej postaci, na przykład w wersji łatwej do czytania lub w innym języku, prosimy powiedzieć nam o tym.**

**Prosimy poinformować nas również, jeżeli potrzebowaliby Państwo usługi tłumaczenia ustnego lub innej pomocy.**

## **Punjabi**

**ਜੇ ਤੁਹਾਨੂੰ ਇਹ ਜਾਣਕਾਰੀ ਕਿਸੇ ਹੋਰ ਰੂਪ ਵਿਚ, ਜਿਵੇਂ ਪੜ੍ਹਨ ਵਿਚ ਆਸਾਨ ਰੂਪ ਜਾਂ ਕਿਸੇ ਦੂਜੀ ਭਾਸ਼ਾ ਵਿਚ, ਚਾਹੀਦੀ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਸਾਨੂੰ ਦੱਸੋ।**

**ਜੇ ਤੁਹਾਨੂੰ ਦੁਭਾਸ਼ੀਏ ਦੀ ਜਾਂ ਸਹਾਇਤਾ ਦੀ ਲੋੜ ਹੈ ਤਾਂ ਕਿਰਪਾ ਕਰਕੇ ਸਾਨੂੰ ਦੱਸੋ।**

## **Romanian**

**Dacă aveți nevoie de informații în alt format, ca de exemplu caractere ușor de citit sau altă limbă, vă rugăm să ne informați.**

**Dacă aveți nevoie de un interpret sau de asistență, vă rugăm să ne informați.**

## **Traditional Chinese**

**如果您需要以其他方式了解信息，如易读或其他语种，请告诉我们。**

**如果您需要口译人员或帮助，请告诉我们。**